

Black hole solutions in four-dimensional topological gravity**S. Mignemi[†]**

Dipartimento di Matematica, Università di Cagliari
viale Merello 92, 09123 Cagliari, Italy

We study spherically symmetric solutions of a four-dimensional theory of gravity with a topological action, which was constructed as a Yang-Mills theory of the Poincaré group and can be considered a generalization to higher dimensions of well-known two-dimensional models. We also discuss the perturbative degrees of freedom and the properties of the theory under conformal transformations.

[†] e-mail: MIGNEMI@CA.INFN.IT

1. Introduction

In recent years, there has been a great interest in the possibility of writing down the theory of gravity as a Yang-Mills theory of the Poincaré or (anti)-de Sitter group, motivated by the fact that in this form the theory resembles the other known fundamental interactions and should be renormalizable [1,2]. This goal has been achieved in two and three dimensions by using actions of topological origin. In the three-dimensional case, one makes use of a Chern-Simons type action [1], while in two dimensions, in order to construct an action, one is forced to introduce a multiplet of auxiliary scalar fields besides the gravitational variables [2]. After a suitable identification of the Yang-Mills fields with the geometric quantities of a riemannian manifold, these actions result to be linear in the curvature tensor of the manifold, and give rise to models which are the closest possible generalization of general relativity to two or three dimensions.

Some time ago, Chamseddine [3] showed that this approach can be generalized to higher dimensions at the cost of introducing actions containing higher powers of the Riemann tensor. In particular, for odd dimensional theories, one generalizes the Chern-Simons action by using higher dimensional Chern-Simons invariants, while in even dimensions, where a Chern-Simons action cannot be defined, one makes recourse to Gauss-Bonnet invariants coupled to auxiliary scalar fields.

These models are expected to be renormalizable, since no potential counterterms exist [3]. Moreover, as in the lower-dimensional cases, different phases are expected to arise, depending on the vacuum expectation values of the fields. In particular, a topological phase with vanishing expectation value for the vielbeins, and some broken phases including the ordinary spacetime of general relativity should exist.

In this paper, we shall be interested in the four-dimensional case, which, if one excludes possible mechanisms of spontaneous compactification, is the physically relevant one. We shall limit ourselves to the study of the "riemannian" phase, corresponding to vanishing torsion and nonvanishing expectation value for the vierbeins. The physics of the four-dimensional models will turn out to be quite different from that of Einstein gravity. In particular, the field equations do not reduce to those of general relativity in any limit. Moreover, one of the properties of the lower-dimensional models which seems to be valid

also in higher dimensions is the fact that these theories do not possess any propagating degrees of freedom around maximally symmetric backgrounds.

In order to investigate more deeply the physical implications of these models, we shall study the spherically symmetric solutions of the field equations in the riemannian sector of the theory. We show that in general the field equations do not determine completely the form of the solutions. We also examine the propagation of small excitations around the relevant backgrounds and discuss the behaviour of the model under conformal transformations.

2.1 Gauge theories of gravity in two dimensions

Before describing the 4-dimensional theory, we briefly review the properties of the 2-dimensional models [2], which have many similar features. In two dimensions, the gauge invariant action can be written

$$I_2 = \int_{M_2} \epsilon_{ABC} \eta^A F^{BC} \quad (2.1)$$

where ϵ_{ABC} is a totally antisymmetric tensor, F^{AB} is the field strength of $SO(1, 2)$ or $ISO(1, 1)$ and η is a triplet of scalars in the fundamental representation of the gauge group. The group indices run from 0 to 2.

The action is anomaly free and finite [2]. Connection with gravity is made through the identification $A^{ab} = \omega^{ab}$, $A^{a2} = e^a$, where ω^{ab} and e^a are the spin connection and vielbeins of the two-dimensional manifold ($a, b = 0, 1$). This yields $F^{ab} = \mathcal{R}^{ab} + \lambda e^a e^b$, $F^{a4} = \mathcal{T}^a$, where $\mathcal{R}^{ab}$ and $\mathcal{T}^a$ are the curvature and the torsion 2-forms on M_2 and $\lambda = 0, \pm 1$ for the Poincaré or (anti-)de Sitter group respectively.

With these identifications, the action can then be written

$$\int_{M_2} \epsilon_{ab} (\eta^2 \mathcal{R}^{ab} + \eta^a \mathcal{T}^b + \lambda \eta^2 e^a e^b) \quad (2.2)$$

The field equations obtained varying with respect to η^a imply the vanishing of the torsion. One can then define as usual the riemannian connection and the metric $g_{mn} = \eta_{ab} e_m^a e_n^b$, in terms of which the action becomes

$$I_2 = \int_{M_2} \sqrt{-g} \eta [R(g_{mn}) + \lambda] \quad (2.3)$$

where R is the Ricci scalar and $\eta \equiv \eta^2$. In dimensions other than two the scalar field η in front of R can be reabsorbed by rescaling the metric g_{mn} . This however is not possible in two dimensions. This is obvious in view of the fact that $\sqrt{-g}R$ is a total derivative in two dimensions and is in fact proportional to the Euler form. A conformal transformation of the metric by a function of the scalar has essentially the effect of adding some η -derivative terms to the action. For $g_{\mu\nu} \rightarrow \eta g_{\mu\nu}$, for example, the action transforms into

$$I_2 = \int_{M_2} \sqrt{-g} \eta [R + 4 \frac{(\nabla \eta)^2}{\eta^2} + \lambda \eta] \quad (2.4)$$

A weak field expansion of the action (2.3) shows that it possesses no propagating degrees of freedom. Nevertheless, it admits black hole solutions which, if $\lambda \neq 0$, in the Schwarzschild gauge read

$$ds^2 = - \left(\frac{\lambda x^2}{2} - c \right) dt^2 + \left(\frac{\lambda x^2}{2} - c \right)^{-1} dx^2 \quad \eta = \lambda x \quad (2.5)$$

The metric is that of de Sitter or anti-de Sitter spacetime depending on the sign of λ . The integration constant c is proportional to the ADM mass of the solution. In particular, in the anti-de Sitter case, if $c > 0$ an event horizon is present at $x = \sqrt{2c/\lambda}$ and the solution can be interpreted as a regular black hole [4].

If $\lambda = 0$, instead, the spacetime is flat and

$$ds^2 = -(ax + b)dt^2 + (ax + b)^{-1}dx^2 \quad \eta = cx + d \quad (2.6)$$

with a , b , c and d integration constants. Also these solutions can be interpreted as black holes [4].

2.2 Gauge theories of gravity in 4 dimensions

In four dimensions, the relevant groups are the Poincaré group $ISO(1,3)$, the de Sitter group $SO(1,4)$ and the anti-de Sitter group $SO(2,3)$. In order to construct the action, one can make use of the totally antisymmetric group invariant tensor ϵ_{ABCDE} , in the case of de Sitter or anti-de Sitter groups, while the Poincaré case is recovered by Inönü-Wigner contraction. The group indices run from 0 to 4. The action is given by [3]:

$$I = \int_{M_4} \epsilon_{ABCDE} \eta^A F^{BC} F^{DE} \quad (2.7)$$

where η^A is a multiplet of scalars in the fundamental representation of the gauge group and the field strength 2-form F^{AB} is defined as $F^{AB} = dA^{AB} + A^{AC}A^{CB}$, the 1-form A^{AB} being the connection of the gauge group.

In analogy with the two-dimensional case, one can then identify $A^{ab} = \omega^{ab}$, $A^{a4} = e^a$, where ω^{ab} and e^a are the spin connection and vielbeins of the four-dimensional manifold. This implies that $F^{ab} = \mathcal{R}^{ab} + \lambda e^a e^b$, $F^{a4} = \mathcal{T}^a$, where $\mathcal{R}^{ab}$ and $\mathcal{T}^a$ are the curvature and the torsion 2-forms of the 4-dimensional manifold and $\lambda = 1$ for $SO(2, 3)$, $\lambda = -1$ for $SO(1, 4)$ or $\lambda = 0$ for $ISO(1, 3)$. The action can be written in terms of the geometrical quantities as

$$I = \int_{M_4} \epsilon_{abcd} [\eta^4 (\mathcal{R}^{ab} + \lambda e^a e^b) (\mathcal{R}^{cd} + \lambda e^c e^d) + 4\eta^a \mathcal{T}^b \mathcal{R}^{cd}] \quad (2.8)$$

The field equations obtained by varying (2.8) with respect to the η fields are thus

$$\begin{aligned} \epsilon_{abcd} \eta^4 (\mathcal{R}^{ab} + \lambda e^a e^b) (\mathcal{R}^{cd} + \lambda e^c e^d) &= 0 \\ \epsilon_{abcd} \mathcal{T}^b (\mathcal{R}^{cd} + \lambda e^c e^d) &= 0 \end{aligned} \quad (2.9)$$

It is easy to see that the second equation is satisfied if the torsion vanishes. Contrary to the 2-dimensional case, however, this is not the unique solution. Nevertheless, in the following we shall limit our attention only to this sector.

When the torsion vanishes, the connection can be expressed as usual in terms of the vielbeins, and then one can write the action in the metric formalism as

$$I = \int_{M_4} \sqrt{-g} d^4x \eta (S + 4\lambda R + 24\lambda^2) \quad (2.10)$$

where η is the scalar field η^4 , R is the Ricci scalar on M_4 , and S is the Gauss-Bonnet term

$$S = R_{mnpq} R_{mnpq} - 4R_{mn} R_{mn} + R^2 \quad (2.11)$$

which is proportional to the Euler density in four dimensions. $\sqrt{-g}S$ is therefore a total derivative in 4 dimensions and so can appear in the action only coupled to a scalar. This term has already been introduced in several contexts, due to its peculiar properties. In particular, it is the only four-dimensional higher-derivative term that does not involve

derivatives higher than second order in the field equations [5]. Moreover it does not introduce new degrees of freedom into the action, besides the graviton [6]. For this reasons, it has been considered in Kaluza-Klein models [7] and appears in low-energy string effective actions [8,9].

3.1. The Poincaré group

When $\lambda = 0$, only the first term in (2.10) survives. The action is therefore a total derivative times a scalar and does not include terms proportional to the Einstein-Hilbert action. In the following, in analogy to some 2-dimensional models, we shall also consider the slightly more general case in which a cosmological constant term $\Lambda\eta$ is added to the action:

$$I = \int \sqrt{-g} d^4x \eta(S + \Lambda) \quad (3.1)$$

Of course, if $\Lambda \neq 0$, the action is not gauge-invariant. The field equations arising from the action (3.1) are:

$$H_{mn} = \frac{\Lambda}{2} \eta g_{mn} \quad (3.2)$$

$$S = \Lambda \quad (3.3)$$

where

$$\begin{aligned} H_{mn} = & 2[4R_{p(m}\nabla_{n)}\nabla_p\eta - 2R_{mn}\nabla^2\eta - 2g_{mn}R_{pq}\nabla_p\nabla_q\eta \\ & - 4R\nabla_{(m}\nabla_{n)}\eta + g_{mn}R\nabla^2\eta + 2R_{qmpn}\nabla_p\nabla_q\eta] \end{aligned} \quad (3.4)$$

If we denote by H the trace of H_{mn} :

$$H = 2(2R_{pq}\nabla_p\nabla_q\eta - R\nabla^2\eta) = 4G_{pq}\nabla_p\nabla_q\eta \quad (3.5)$$

the trace of (3.2) gives:

$$H = \Lambda\eta \quad (3.6)$$

The tensor H_{mn} satisfies the equivalent of a Bianchi identity:

$$\nabla_m H_{mn} = \frac{1}{2} \nabla_n H$$

From (3.6) then follows that

$$\nabla_m H_{mn} = \frac{\Lambda}{2} \nabla_n \eta \quad (3.7)$$

consistently with (3.2).

In the following we shall be mainly interested in spherically symmetric solutions of the field equations. If one assumes for the metric the static Schwarzschild form

$$ds^2 = -f^2(r)dt^2 + h^{-2}(r)dr^2 + r^2d\Omega^2 \quad (3.8)$$

the field equations (2) yield:

$$\begin{aligned} (1 - h^2)\eta'' + \left(\frac{h'}{h} - 3hh'\right)\eta' &= -\frac{\Lambda}{4}r^2\eta \\ (1 - 3h^2)\frac{f'}{f}\eta' &= -\frac{\Lambda}{4}r^2\eta \\ h^2f'\eta'' + \left(\frac{h'}{h}f' + h^2f''\right)\eta' &= \frac{\Lambda}{4}r\eta \end{aligned} \quad (3.9)$$

Due to the identity (3.7), only two of these equations are independent. Moreover, recalling that $\sqrt{-g}S$ is a total derivative, one can write (3.3) as:

$$8[f'h(1 - h^2)]' = -\Lambda fh^{-1}r^2 \quad (3.10)$$

3.2 $\Lambda = 0$

In the case of vanishing cosmological constant Λ , a trivial solution of the field equations is given by flat space. In this case η remains undetermined, since H_{mn} vanishes independently of η . The possibility of choosing freely one arbitrary function in the solution seems to be a general feature of the models we are considering. All solutions we were able to find depend on one arbitrary function and only by imposing further constraints can one obtain solutions in closed form.

The flat space solution with constant η can be considered as the ground state of the theory. Let us now look for more general solutions of the field equations. When $\Lambda = 0$, one has from (3.10)

$$f'h(1 - h^2) = \text{const} = \alpha \quad (3.11)$$

For $\alpha = 0$, two possible solutions emerge: either $h = 1$ or f is a constant, which can be put to 1 by rescaling t .

Let us consider the first case: the equations (3.9) are then satisfied if $f'\eta' = 0$. Hence one recovers the flat solution if $f' = 0$, or a solution with arbitrary f and constant η . The second case corresponds to a spacetime with flat spatial sections, and metric of the form

$$ds^2 = -f^2(r)dt^2 + dr^2 + r^2d\Omega^2 \quad (3.12)$$

The arbitrariness of f does not consent in this case to draw conclusions about the causal structure of the solutions. Also the newtonian limit is undetermined, since the newtonian potential is related to f .

In the case $f = 1$, the equations (3.9) reduce to

$$(1 - h^2)\eta'' + \left(\frac{h'}{h} - 3hh'\right)\eta' = 0 \quad (3.13)$$

This equation admits solutions with arbitrary h and η either constant or given by

$$\eta' = \frac{C}{h(1 - h^2)} \quad (3.14)$$

with C an integration constant. The metric has in this case the form

$$ds^2 = -dt^2 + \frac{dr^2}{h^2(r)} + r^2d\Omega^2 \quad (3.15)$$

and is therefore the direct product of the time coordinate with an arbitrary spherically symmetric 3-space. Again, the properties of the specific solutions depend on the form of the function h . However, from the geodesic equation it is easy to see that a point particle does not experience any gravitational force in this metric.

We pass now to consider the case in which $\alpha \neq 0$. In this case one cannot find the general solution of the field equations. Some special solutions can nevertheless be obtained by imposing suitable ansätze. For example, a solution is easily obtained by putting $h^2 = 1/3$, in order to satisfy the second of equations (3.9). The other field equations then imply that $f = ar$ and $\eta = br + c$, with a, b, c , integration constants. This solution is regular everywhere, except for a conical singularity at the origin.

Another class of solutions is obtained for $\eta = \text{const}$. In this case, equations (3.9) are trivially satisfied and from (3.11) one has

$$f' = \frac{\alpha}{h(1 - h^2)} \quad (3.16)$$

with arbitrary h . A special solution can be obtained if one requires $f=h$, as for the Schwarzschild metric of general relativity. In this case, (3.11) can be written as

$$[(1 - h^2)^2]' = a^2 \quad (3.17)$$

which can then be easily integrated yielding

$$h^2 = f^2 = 1 \pm a\sqrt{r-b} \quad (3.18)$$

with a, b integration constant. The curvature invariants built from this metric are singular at $r = 0$ and $r = b$. This indicates the presence of a physical singularity at these points. The singularity at $r = b + a^{-2}$ arising when the minus sign is chosen in (3.18), is instead simply a coordinate singularity, corresponding to a horizon. It is easy to check that in any case, however, the singularities at $r = 0$ or $r = b$ are naked, so that no regular black hole solution exists of this form. It may be interesting to notice, however, that this solution corresponds to a gravitational force decreasing as $r^{-1/2}$ at infinity in the weak field limit.

3.3 $\Lambda \neq 0$

In the case $\Lambda \neq 0$, maximally symmetric solutions exist if $\Lambda > 0$ and are given by de Sitter or anti-de Sitter space with curvature $\pm\sqrt{\Lambda}/24$. In this case, the scalar field η must vanish according to the field equations. This solution can be considered the ground state of the theory.

More general solutions are much more difficult to obtain than in the $\Lambda = 0$ case, since now (3.10) cannot be integrated explicitly. One can however obtain a first integral of (3.10) by writing the metric as:

$$ds^2 = -Adt^2 + \frac{dr^2}{AR^4} + R^2d\Omega^2 \quad (3.19)$$

with $A = A(r)$, $R = R(r)$. In this case,

$$A'R^2(1 - AR^4R'^2) = 2\Lambda r + a \quad (3.20)$$

and for any given R one can obtain A by solving a non-linear first order differential equation. In general it is not possible to give the solution in a closed form. A special (unphysical) case is given by $R = r^{-1/4}$, $A = \Lambda r^{5/2}$, which in different coordinates can be written

as

$$ds^2 = -\Lambda\rho^{5/3}dt^2 + (\Lambda\rho^{5/3})^{-1}d\rho^2 + \rho^{-3}d\Omega^2 \quad (3.21)$$

An explicit solution can also be obtained, as in the $\Lambda = 0$ case, by imposing the ansatz $f = h$. One has now from (3.10):

$$[(1 - h^2)^2]' = \frac{\Lambda r^3}{6} + a \quad (3.22)$$

and then

$$h^2 = f^2 = 1 \pm \sqrt{\frac{\Lambda r^4}{24} + ar + b} \quad (3.23)$$

with a and b integration constants. With this form of the metric functions, the field equations (3.9) can be satisfied only if $\eta = 0$. If $a = b = 0$, the metric reduces to that of anti-de Sitter or de Sitter spacetime, depending on the sign in (3.23). Otherwise, the solution is de Sitter or anti-de Sitter only asymptotically. The curvature invariants are singular at $r = 0$ and at the zeroes of $\frac{\Lambda r^4}{24} + ar + b$. In general also some horizons may be present, depending on the signs of the parameters of the solutions. However, the only case in which no naked singularity arise is when the minus sign is taken in (3.23) and $\Lambda > 0$, $a < 0$, $b > \max\{1, 2\Lambda^{-1/3}(3a/4)^{4/3}\}$. In this case the metric is qualitatively similar to that of a black hole in de Sitter spacetime.

4. The (anti)-de Sitter group

In the (anti)-de Sitter case, the action takes the form:

$$I = \int \sqrt{-g} d^4x \eta(S + 4\lambda R + 24\lambda^2) \quad (4.1)$$

and the field equations are

$$H_{mn} + 4\lambda\eta G_{mn} + 4\lambda K_{mn} - 12\lambda^2\eta g_{mn} = 0 \quad (4.2)$$

$$S + 4\lambda R + 24\lambda^2 = 0 \quad (4.3)$$

where G_{mn} is the Einstein tensor and

$$K_{mn} = -(\nabla_m \nabla_n - g_{mn} \nabla^2)\eta \quad (4.4)$$

Even if a Einstein-Hilbert term is now present in the action, the theory is quite different from general relativity, since the coupling with the scalar field η yields the constraint (4.3), which has no counterpart in the Einstein theory.

It is difficult to find a general spherically symmetric solution of the field equations. However, from (2.9) one sees that the field equations can be satisfied if $\mathcal{R}^{23} = -\lambda e^2 e^3$. Substituting the ansatz (3.8) one can then check that any metric of the form

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{1 + \lambda r^2} + r^2 d\Omega^2 \quad (4.5)$$

with arbitrary f is a solution of the field equations. These solutions are therefore a direct generalization of the solutions (3.12), which correspond to the $\lambda = 0$ limit. For $\lambda < 0$, they describe a spacetime whose spatial sections are 3-spheres. In particular, in the special case $f(r) = h(r) = 1 + \lambda r^2$, one obtains the de Sitter solution. In the general case, instead, the properties of the solutions depend on the specific form of the function f . Analogous considerations hold for $\lambda > 0$.

For the solutions (4.5), the scalar field equations (4.2) become, if $f \neq h$,

$$\frac{\eta'}{r} = \lambda \eta \quad \eta'' + \frac{\lambda r \eta'}{1 + \lambda r^2} = \lambda \eta \quad (4.6)$$

The only solution of this system is $\eta = 0$. If $f = h$, instead, all equations are identically satisfied and η can be an arbitrary function of r .

We point out that consideration of the ansatz $\mathcal{R}^{01} = -\lambda e^0 e^1$ does not generate any new solutions.

5. Conformal transformations

In this section, we briefly discuss the effect of a conformal transformation on the action (2.10). In four dimensions, a suitable conformal transformation allows one to remove the scalar field in front of the Einstein-Hilbert term, but the effect on the Gauss-Bonnet term is simply to add some term containing fourth order derivatives of the scalar fields to the action. This is analogous to what happens in two dimensions with the Einstein-Hilbert term, as discussed in section 2.1. The transformation of course affects the geometrical properties of the theory, but not the field theoretical ones, since it simply amounts to a

redefinition of the fields. In particular, the field equations will contain at most second order derivatives of the fields, even if not linearly.

In more detail, if we consider the action (2.10) and perform the conformal transformation $g_{mn} \rightarrow e^{2\phi} g_{mn}$, where $\eta = e^{2\phi}$, in order to eliminate the scalar field in front of R , the action becomes:

$$I = 4\lambda \int \sqrt{-g} d^4x \{ R + 6(\nabla\phi)^2 + 6\lambda e^{2\phi} + \frac{1}{4\lambda} e^{-2\phi} [S + 2R(\nabla\phi)^2 - 2R_{ab}\nabla_a\phi\nabla_b\phi - 8(\nabla^2\phi)^2 + 32\nabla^2\phi(\nabla\phi)^2] \} \quad (5.1)$$

Hence, in the $\lambda \neq 0$ case, the rescaled action assumes the usual Einstein-Hilbert form with some non-minimal corrections and displays some similarity with the effective four-dimensional string lagrangians [9]. One might speculate that effective string lagrangians can be obtained in this way from a suitable modification of the gravitational gauge group, as in two dimensions [10]. When $\lambda = 0$, instead, only the terms in square brackets survive and the action changes only by higher derivative terms in the scalars. It may be interesting to notice that the peculiar combination of these scalar terms does not lead to derivatives higher than second in the field equations.

In the Poincaré case, solutions of (5.1) can be easily obtained from those of the conformally equivalent action by simply applying the conformal transformation to the metric. In the (anti-)de Sitter case, instead, this is not possible, since we were only able to find solutions with vanishing scalar field, for which the conformal transformation becomes singular.

6. Linearization

In order to investigate the particle content of the models discussed above, it is interesting to consider the propagation of excitations around given backgrounds. It is well known that the results depend on the specific ground state around which one expands: for example, in the unbroken phase of the full theory, $e = \omega = 0$, no propagation can arise. Here we consider the theory in its metric form (2.10). As we shall see, also in this case no propagating degree of freedom is present, at least for the most natural choice of the ground state. This is similar to the two-dimensional case, where it is well-known that no propagation takes place.

In order to investigate this point, one must expand the metric and the scalar around a fixed ground state, $g_{mn}^{(0)}$, $\eta^{(0)}$, as

$$g_{mn} = g_{mn}^{(0)} + h_{mn}, \quad \eta = \eta^{(0)} + \phi$$

For generic background the linearized action has a complicated form. However, for maximally symmetric backgrounds, one has in four dimensions (the superscripts denote the order of the terms in the expansion):

$$\begin{aligned} (\sqrt{-g}R)^{(1)} &= -(\nabla^2 h_{aa} - \nabla_a \nabla_b h_{ab}) - 3\lambda h_{aa} \\ (\sqrt{-g}S)^{(1)} &= 4\lambda(\nabla^2 h_{aa} - \nabla_a \nabla_b h_{ab}) \\ (\sqrt{-g}R)^{(2)} &= -\frac{1}{4}(h_{ab}\nabla^2 h_{ab} - h_{aa}\nabla^2 h_{bb} + 2h_{cc}\nabla_a \nabla_b h_{ab} - 2h_{ac}\nabla_a \nabla_b h_{bc}) \\ &\quad - \lambda(2h_{ab}h_{ab} - \lambda h_{aa}h_{bb}) \\ (\sqrt{-g}S)^{(2)} &= 0 \end{aligned}$$

where $(\sqrt{-g}S)^{(2)}$ vanishes in four dimensions because of the topological properties of the Gauss-Bonnet term [6].

For $\lambda = 0$, the most natural choice for the background is given by the flat Minkowski metric with constant scalar. It follows that $(\sqrt{-g}\eta S)^{(2)} \sim (\sqrt{-g}S)^{(1)}\eta^{(1)} = 0$, since $(\sqrt{-g}S)^{(1)} = 0$ if $\lambda = 0$. Analogously, for $\lambda \neq 0$, defining $\mathcal{L} = S + 4\lambda R + 24\lambda^2$,

$$\begin{aligned} \sqrt{-g}\mathcal{L}^{(1)} &= 0 \\ (\sqrt{-g}\mathcal{L})^{(2)} &= -\frac{1}{4}(h_{ab}\nabla^2 h_{ab} - h_{aa}\nabla^2 h_{bb} + 2h_{cc}\nabla_a \nabla_b h_{ab} - 2h_{ac}\nabla_a \nabla_b h_{bc}) \end{aligned}$$

From the results of section 4, the most natural choice for the ground state appears to be $\eta^{(0)} = 0$. Hence

$$(\sqrt{-g}\eta\mathcal{L})^{(2)} = \eta^{(1)}(\sqrt{-g}\mathcal{L})^{(1)} + \eta^{(0)}(\sqrt{-g}\mathcal{L})^{(2)} = 0$$

Therefore, for any value of λ no propagating degree of freedom is present in the spectrum of the theory, since the term of the lagrangian quadratic in the perturbations vanishes. However, if one had expanded around $\eta = \text{const}$ in the de Sitter case, a graviton would have appeared in the spectrum, since in that case $(\sqrt{-g}\eta\mathcal{L})^{(2)} \sim (\sqrt{-g}\mathcal{L})^{(2)}$ which is proportional to the graviton propagator.

Also interesting is the case of the non-gauge-invariant action (3.1) with $\Lambda \neq 0$. In this case, taking an (anti-)de Sitter background with $\eta^{(0)} = 0$,

$$(\sqrt{-g}\eta(S + \Lambda))^{(2)} = \eta^{(1)}(\sqrt{-g}(S + \Lambda))^{(1)} = \phi(\nabla^2 g_{ab} - \nabla_a \nabla_b)h_{ab} + \frac{\Lambda}{2}\phi h_{aa}$$

In the gauge $\nabla_b h_{ab} = \frac{1}{2}\nabla_b h_{aa}$, the propagator reduces to $\frac{1}{2}\phi(\nabla^2 + \Lambda)h$, with $h = h_{aa}$, or diagonalizing

$$\frac{1}{2}[(\phi - h)(\nabla^2 + \Lambda)(\phi - h)] - \frac{1}{2}[(\phi + h)(\nabla^2 + \Lambda)(\phi + h)]$$

This corresponds to two propagating massive scalar degrees of freedom, one of which is a ghost.

7. Final remarks

We have discussed some properties of a Yang-Mills theory of gravity in four dimensions. From the results obtained, it appears that the field equations for the metric are somehow underdetermined, at least in the case of a spherically symmetric ansatz. Only one of the metric fields is in fact determined by the field equations. On the contrary, it is quite difficult to find non-trivial solutions for the scalar field. In particular, in the de Sitter and anti-de Sitter cases, all the solutions we have found imply vanishing scalar field. A consequence of this fact is that no propagating degrees of freedom are present in the spectrum of the theory. This was perhaps to be expected in view of the topological nature of the action of the theory.

Of course, as in all gravitational theories, it is essential for the physical interpretation to consider the coupling to matter. If one assumes that the scalar field η is physical, and not simply a Lagrange multiplier, the coupling is not uniquely determined. This is a common feature in gravity-scalar theories. However, even introducing non-trivial couplings of the matter with the scalar field, it does not seem possible to obtain a newtonian limit in the framework of these models. A more physical behaviour may perhaps be obtained by considering conformally related models, as those discussed in section 4, in which an Einstein-Hilbert term is present in the action for $\lambda \neq 0$.

Some of the problems discussed so far might be solved by considering the more general sector containing non-trivial torsion fields, which could give rise to more general solutions of the field equations.

References

- [1] E. Witten, Nucl. Phys. **B311**, 46 (1989);
- [2] T. Fukuyama and K. Kamimura, Phys. Lett. **B160**, 259 (1985); K. Isler and C. Trugenberger, Phys. Rev. Lett. **63**, 834 (1989); A.H. Chamseddine and D. Wyler, Phys. Lett. **B228**, 75 (1989);
- [3] A.H. Chamseddine, Nucl. Phys. **B346**, 213 (1990);
- [4] M. Cadoni and S. Mignemi, Phys. Rev. **D51**, 4319 (1995); Phys. Lett. **B358**, 217 (1995);
- [5] D. Lovelock, J. Math. Phys. **12**, 498 (1971);
- [6] D.G. Boulware and S. Deser, Phys. Rev. Lett. **55**, 2656 (1985);
- [7] J. Madore, Phys. Lett. **A110**, 289 (1985); F. Müller-Hoissen, Phys. Lett. **B163**, 106 (1985);
- [8] B. Zwiebach, Phys. Lett. **B156**, 315 (1985);
- [9] D.J. Gross and J.H. Sloan, Nucl. Phys. **B291**, 41 (1987);
- [10] H. Verlinde, in *Sixth Marcel Grossmann Meeting on General Relativity*, M. Sato and T. Nakamura, eds. (World Scientific, 1992); D. Cangemi and R. Jackiw, Phys. Rev. Lett. **69**, 233 (1992).